



Heart Attack



Disclosures

- Relationship with Novartis Pharmaceuticals Corporation, Amgen, Amarin, Bayer, Pfizer, Lexicon Pharmaceuticals, and Idorsia that includes consulting or advising.
- Relationship with Janssen that includes research grant funding paid directly to the research department

Objectives



We will

- review the continuum of coronary artery disease from acute coronary syndrome to chronic management.
- Discuss chronic management through lifestyle and the importance of pharmacologic treatment

Not all MI's are the same: Type 1 - 5

TABLE A Universal Classification of MI

Type 1: Spontaneous MI

Spontaneous MI related to atherosclerotic plaque rupture, ulceration, fissuring, erosion, or dissection with resulting intraluminal thrombus in ≥ 1 of the coronary arteries leading to decreased myocardial blood flow or distal platelet emboli with ensuing myocyte necrosis. The patient may have underlying severe CAD, but on occasion nonobstructive or no CAD.

Type 2: MI secondary to ischemic imbalance

In instances of myocardial injury with necrosis where a condition other than CAD contributes to an imbalance between MVO₂, e.g., coronary endothelial dysfunction, coronary artery spasm, coronary embolism, tachy-/bradyarrhythmias, anemia, respiratory failure, hypotension, and hypertension with or without LVH.

Type 3: MI resulting in death when biomarker values are unavailable

Cardiac death with symptoms suggestive of myocardial ischemia and presumed new ischemic changes or new LBBB, but death occurred before blood samples could be obtained, before cardiac biomarker testing, or before ECG testing.



Type 5: MI related to CAD

MI associated with CABG, percentile URL). In addition, (iii) imaging evidence of new loss of

CABG indicates coronary artery bypass grafting; MVO₂, myocardial oxygen consumption; Modified from Thygesen et al.

Not all MI's are the same: Type 1 - 5

TABLE A Universal Classification of MI

Type 1: Spontaneous MI

STEMI or NSTEMI

Spontaneous MI related to atherosclerotic plaque rupture, ulceration, fissuring, erosion, or dissection with resulting intraluminal thrombus in ≥ 1 of the coronary arteries leading to decreased myocardial blood flow or distal platelet emboli with ensuing myocyte necrosis. The patient may have underlying severe CAD, but on occasion nonobstructive or no CAD.

Type 2: MI secondary to ischemic imbalance

Demand Ischemia/infarct NOT STEMI or NSTEMI

In instances of myocardial injury with necrosis where a condition other than CAD contributes to an imbalance between MVO₂, e.g., coronary endothelial dysfunction, coronary artery spasm, coronary embolism, tachy-/bradyarrhythmias, anemia, respiratory failure, hypotension, and hypertension with or without LVH.

Type 3: MI resulting in death when biomarker values are unavailable

Cardiac death with symptoms suggestive of myocardial ischemia and presumed new ischemic electrocardiographic changes or new LBBB, but death occurred before blood samples could be obtained, before cardiac biomarker could rise, or in rare cases where blood was not collected for cardiac biomarker testing.

Type 4a: MI related to PCI

MI associated with PCI is arbitrarily defined by elevation of cTn values $> 5 \times$ 99th percentile URL in patients with normal baseline values (< 99 th percentile URL) or a rise of cTn values $> 20\%$ if baseline values are elevated and are stable or falling. In addition, either (i) symptoms suggestive of myocardial ischemia, (ii) new ischemic electrocardiographic changes or new LBBB, (iii) angiographic loss of patency of a major coronary artery or a side branch or persistent slow or no flow or embolization, or (iv) imaging demonstration of new loss of viable myocardium or new regional wall motion abnormality is required.

Type 4b: MI related to stent thrombosis

MI associated with stent thrombosis is detected by coronary angiography or autopsy in the setting of myocardial ischemia and with a rise and/or fall of cardiac biomarker values with ≥ 1 value above the 99th percentile URL.

Type 5: MI related to CABG


MI associated with CABG is arbitrarily defined by elevation of cardiac biomarker values $> 10 \times$ 99th percentile URL in patients with normal baseline cTn values (< 99 th percentile URL). In addition, either (i) new pathological Q waves or new LBBB, or (ii) angiographically documented new graft or new native coronary artery occlusion, or (iii) imaging evidence of new loss of viable myocardium or new regional wall motion abnormality.

CABG indicates coronary artery bypass graft; CAD, coronary artery disease; cTn, cardiac troponin; LBBB, left bundle-branch block; LVH, left ventricular hypertrophy; MI, myocardial infarction; MVO₂, myocardial oxygen consumption; PCI, percutaneous coronary intervention; and URL, upper reference limit.


Modified from Thygesen et al. (21).

Acute Coronary Syndrome/Chronic Stable


Stable Angina



NSTE-ACS



STEMI




Acute Tools of the “trade”



<https://www.plumbing-draincleaning.com/drain-cleaning.html>




www.plumbingsupply.com/%2Fflogbusters.html&psig=AOvVaw3iT0dXRoxDvHDBT5VcXh4y&ust=1668264017895000&source=images&cd=vfe&ved=0CBEQ3YkBahcKEwjYtqeVr6b7AhUAAAAAHQAAAwAAQCz





www.amazon.com/%2FUpgraded-Anti-break-Plumbing-Bathroom-Cleaning%2Fdp%2F809GK99MQ4&psig=AOvVaw3iT0dXRoxDvHDBT5VcXh4y&ust=1668264017895000&source=images&cd=vfe&ved=0CA0Q3YkBahcKEwjYtqeVr6b7AhUAAAAAHQAAAwAw


Tools of the “trade”



Acute Tools of the “trade”



Coronary Artery Bypass Graft (CABG)




Coronary Calcium and statin eligibility (2019 GL)




CAC = 0 → NO STATIN

CAC = 1-99 → Favor
STATIN

CAC = ≥ 100 or $\geq 75^{\text{th}}$
Percentile → **STATIN**



**POPULATION:**

Framingham (Offspring and 3rd Generation).
 50±10 yrs of age. Female 50.9%.

MAJOR CVD included:

1 coronary heart disease (CHD),
 2 stroke, and
 3 peripheral arterial disease.

Additionally, authors included

4 MI, and


5 death from CHD (i.e., fatal coronary event, MI, or cerebrovascular accident [i.e., ischemic stroke, hemorrhagic stroke]).

Right Arm:
Systolic Pressure mmHg

Right Ankle:
Systolic Pressure
Posterior Tibial (PT) mmHg
Dorsalis Pedis (DP) mmHg

Left Arm:
Systolic Pressure mmHg

Left Ankle:
Systolic Pressure
Posterior Tibial (PT) mmHg
Dorsalis Pedis (DP) mmHg



Ankle-Brachial Index Interpretation

Above 0.90: Normal
0.71 - 0.90: Mild Obstruction
0.41 - 0.70: Moderate Obstruction
0.00 - 0.40: Severe Obstruction

Right ABI equals Ratio of:

Higher of the Right Ankle Pressures (PT or DP)
Higher Arm Pressure (right or left arm)

$$\frac{\frac{\text{mmHg}}{\text{mmHg}}}{\text{mmHg}} = \square \cdot \square \square^*$$

Left ABI equals Ratio of:

Higher of the Left Ankle Pressures (PT or DP)
Higher Arm Pressure (right or left arm)


$$\frac{\frac{\text{mmHg}}{\text{mmHg}}}{\text{mmHg}} = \square \cdot \square \square^*$$

* The lower of these numbers is the patient's overall ABI.


Overall ABI (lower ABI) = _____

Vessel Disease


Calcified Vessel



Normal



Mild PAD



Moderate PAD



Severe PAD



ABI

> 1.4

0.9 - 1.4

0.7 - 0.89

0.51 - 0.69

≤ 0.5

TBI

unaffected






> 0.6

0.34 - 0.59






0.12 - 0.34

≤ 0.11



Doppler



PVR



5-Year KM Estimates of ACM and Severe Vascular Events^{5†}



Persons at risk

	0	1	2	3	4	5	6	7	8	9	10
PAD no/unknown	5392	5303	5192	5085	5017	4935	4464	4004	3953	3885	3850 [‡]
Asymptomatic PAD	836	810	776	742	722	700	612	528	520	493	484 [‡]
Symptomatic PAD	593	561	515	484	463	433	357	301	286	276	268 [‡]

Older: 72
 Female: 58%
 ABI >1.5 excluded


OUTCOMES:

1 all-cause mortality
 OR severe vascular events
 2 myocardial infarction,
 3 coronary revascularization,
 4 stroke,
 5 carotid revascularization,
 6 peripheral revascularization, or
 7 amputation


Older: 72
 Female: 58%
 ABI >1.5 excluded

OUTCOMES:


1 all-cause mortality
 OR severe vascular
 events
 2 myocardial
 infarction,
 3 coronary
 revascularization,
 4 stroke,
 5 carotid
 revascularization,
 6 peripheral
 revascularization, or
 7 amputation



Where would you find a stand of trees that would most likely yield apples?



https://web.extension.illinois.edu/apples/images/us_map.gif




https://en.wikipedia.org/wiki/Climate_of_the_United_States

Which stand of trees would you most likely find apples?




Heart Disease Death Rates, 2014-2016 Adults, Ages 35 +, by County




Heart Disease Death Rates, 2014-2016 Adults, Ages 35 +, by County

@VietHeartPA




<https://www.cdc.gov/heartdisease/facts.html>

Self-reported High Total Cholesterol Among Adults, 2017*




<https://www.cdc.gov/cholesterol/facts.html>

*Data Source: BRFSS, Adults (20+) who answered "yes" to the question, "Have you ever been told by a doctor, nurse or other health professional that your blood cholesterol is high?"



SMOKING

<https://www.cdc.gov/vitalsigns/tobaccouse/smoking/infographic.html>




<https://www.cdc.gov/diabetes/data/center/slides.html>

Traditional risk factors in First MI

WAIT!!

- A significant number of folks with 1st MI also have 0 RF; in addition, they may have an increased risk of death.
- In 542,008 patients presenting with a first myocardial infarction: the percentage with **0, 1, 2, 3, and 4 risk factors was 14.4%, 34.1%, 31.6%, 15.4%, and 4.1%**, respectively

Risk Factors:
Hypertension
Smoking
Dyslipidemia
Diabetes
Family Hx of CAD



SMuRF-Less

Intermountain data presented at ACC 22. Patients with 1st STEMI from 2000-2021 comparing those with **standard modifiable risk factors (SMuRF)*** and those without **SMuRF-Less**.

- STEMI pts (n=3,510), SMuRF-Less made up over 1 in 4 pts, or 26.2% (n=919).
- SMuRF-Less pts were younger, more frequently male, and had fewer overall co-morbidities
- While unadjusted HR for MACE favored SMuRF-Less, an adjusted HR demonstrated similar outcomes other than persistent lower HF admissions.

*Diabetes, hypertension, smoking, hyperlipidemia



Journal of
Clinical Medicine

an Open Access Journal by MDPI

Cardiovascular Outcomes of ST-Elevation Myocardial Infarction (STEMI) Patients without Standard Modifiable Risk Factors (SMuRF-Less): The Intermountain Healthcare Experience

Jeffrey L. Anderson; Stacey Knight; Heidi T. May; Viet T. Le; Jawad Almajed; Tami L. Bair; Kirk U. Knowlton; Joseph B. Muhlestein

J. Clin. Med. **2023**, Volume 12, Issue 1, 75

@VietHeartPA




A. Demographics	SMuRF		SMuRF-less	
	n=2591		n=919	
	n	%	n	%
Age groups				
<40	85	3.28%	49	5.33%
40-49	360	13.89%	140	15.23%
50-59	720	27.79%	228	24.81%
60-69	717	27.67%	271	29.49%
70-79	471	18.18%	150	16.32%
>79	238	9.19%	80	8.71%
Gender				
Male	1885	72.75%	709	77.15%
Female	706	27.25%	210	22.85%
Race				
White/Caucasian	2260	87.23%	818	89.01%
African American	14	0.54%	8	0.87%
Asian	57	2.20%	15	1.63%
Pacific Islander	5	0.19%	3	0.33%
Unknown	255	9.84%	75	8.16%


You have a patient with Atherosclerosis. Now WHAT?

@VietHeartPA


Stable Angina/Claudication




NSTE-ACS/Acute limb ischemia



STEMI/Stroke/Amputation



Secondary Prevention: Avoiding a 2nd Event



Find the culprits for future problems

Risk Factors

- Hypertension
- Smoking
- Dyslipidemia
- Diabetes
- Family history of CAD



<https://www.tandfonline.com/doi/10.1080/1372193/2017.1372193>
 F2092098-Know-Your-Numbers&psig=AOvVaw1hJasK6jFWqkMS4GtzZaTK&ust=1668266362677000&source=images&cd=vfe&ved=0CBEQ3YkBaHcKEwig2r7Ttqb7AhUAAAAAHQAAAAQCA




www.tandfonline.com/doi/10.1080/1372193/2017.1372193
 oi%2Fpdf%2F10.1080%2F14779072.2017.1372193&psig=AOvVaw3LYXMOz7MMRgNzhUxbxRui&ust=1668266458800000&source=images&cd=vfe&ved=0CBEQ3YkBaHcKEwiguKSQt6b7AhUAAAAHQAQAAAAQDA



<https://www.tandfonline.com/doi/10.1080/1372193/2017.1372193>
 smokers&psig=AOvVaw2SpAccR8J5kzAqJdyxMij1&ust=1668265641971000&source=images&cd=vfe&ved=0CBAQ3YkBaHcKEwigQmMv5s6b7AhUAAAAAHQAAAAQBA




www.genengnews.com/2017/07/20/new-diabetes-therapy-might-be-found-in-protein-commonly-found-throughout-the-body/
 body%2F&psig=AOvVaw35kYHy3dHbnP8eRYj5AGmt&ust=1668266607632000&source=images&cd=vfe&ved=0CBEQ3YkBaHcKEwiw_Zbwt6b7AhUAAAAAHQAAAAQAw





<https://healthblog.uofmhealth.org/heart-health/what-you-should-know-about-counseling-and-testing-for-genetic-heart-disease>

Antiplatelet(s):

Plaque presence = potential for rupture or thrombus;



<https://giphy.com/gifs/search/myocardial>



1. **Aspirin** 81 mg or 325 mg

- ADAPTABLE trial = either; 81 mg demonstrates same benefit, less bleeding

2. **P2y12 inhibitors**: Clopidogrel 75 mg, Prasugrel 10 mg, or Ticagrelor (90 mg po bid or 60 mg po bid).

3. **Dual antiplatelet therapy (DAPT)**: Both ASA + P2y12i

When to go to ASA or P2y12i alone?


FIGURE 1 Master Treatment Algorithm for Duration of P2Y₁₂ Inhibitor Therapy in Patients With CAD Treated With DAPT




FIGURE 1 Master Treatment Algorithm for Duration of P2Y₁₂ Inhibitor Therapy in Patients With CAD Treated With DAPT

FIGURE 1 Master Treatment Algorithm for Duration of P2Y₁₂ Inhibitor Therapy in Patients With CAD Treated With DAPT





Antiplatelet(s):

Plaque presence = potential for rupture or thrombus;



<https://giphy.com/gifs/search/myocardial>



1. **Aspirin** 81 mg or 325 mg OR **P2y12 inhibitors**
Clopidogrel 75 mg, Prasugrel 10 mg, or
Ticagrelor (90 mg po bid or 60 mg po bid).

As a single agent going forward? CAPRIE, 1996 study demonstrated cardiovascular benefit and less bleeding with clopidogrel over aspirin monotherapy.

Host-Exam 2022 affirmed data from CAPRIE trial of P2Y12i over aspirin.


ULTIMATE-DAPT 2024 POST MI Ticagrelor monotherapy vs DAPT after 30 days.
Ticagrelor>DAPT

Guidelines are still geared to ASA 81 mg monotherapy.


What about Atrial Fibrillation and Coronary Artery Disease?

FIGURE 9 Recommended Duration of Antiplatelet Therapy[†]


@VietHeartPA




What about Atrial Fibrillation and Coronary Artery Disease?



What about Atrial Fibrillation and Coronary Artery Disease?




What about Atrial Fibrillation and Coronary Artery Disease?



What about Atrial Fibrillation and Coronary Artery Disease?

FIGURE 9 Recommended Duration of Antiplatelet Therapy†

@VietHeartPA



ACS indicates acute coronary syndrome; ASA, aspirin; CCD, chronic coronary disease; DAPT, dual antiplatelet therapy; DES, drug-eluting stent; DOAC, direct oral anticoagulant; MI, myocardial infarction; OAC, oral anticoagulants; PCI, percutaneous coronary intervention; SAPT, single antiplatelet therapy. *Colors correspond to Class of Recommendation in Table 3. †This figure does not encompass all recommendations within this section.

Case

55-year-old man returns for annual follow-up.

PMHx: Had an MI at age 50, 2vCABG. Has Paroxysmal Afib.

FMHx: Mom had MI at age 55. Has one sister, A&W.

SocHx: Florist. Single. Lifetime non-smoker, drinks 1-2 beers on the weekends. Lifts weights 2-3 times a week at the gym.

MEDS: Clopidogrel 75 mg, rosuvastatin 40 mg, ezetimibe 10 mg, bi-weekly Repatha 140 mg/mL SC, metoprolol succinate 50 mg. SL NTG 0.4 mg PRN.

Vitals: BP 120/80, HR 55, SaO2 95%, T 98.7, Wt 200 Ht 5'9" BMI 29.5

LABS: TC 200, Trig 110, HDL 42, LDL 50. A1c 5.5%, Fasting Glucose 92 mg/dL

What are your recommendations?

Paroxysmal AF. Antithrombotic regimen?

1. Lifestyle modifications for health
2. Initiate oral anticoagulant and stop P2y12 inhibitor.
3. Watch for bleeding complications of bleeding (e.g., GI)

What about B-blockers in those who have had an acute myocardial infarction

2021 – recommendation in those prescribed beta-blockers during an AMI event, to stop after 3 years if LV Function was “normal” – LVEF $\geq 50\%$

2023 – reassess beta-blocker therapy when LVEF $\geq 50\%$ at >1 year.

2024 – what about, right “out of the gate” with AMI and LVEF $\geq 50\%$?

4.3.2. Beta Blockers

Recommendations for Beta Blockers

Referenced studies that support the recommendations are summarized in the [Online Data Supplement](#).


COR	LOE	Recommendations
1	A	1. In patients with CCD and LVEF $\leq 40\%$ with or without previous MI, the use of beta-blocker therapy is recommended to reduce the risk of future MACE, including cardiovascular death. ¹⁻³
1	A	2. In patients with CCD and LVEF $< 50\%$, the use of sustained release metoprolol succinate, carvedilol, or bisoprolol with titration to target doses is recommended in preference to other beta blockers.* ^{1,3-8}
2b	B-NR	3. In patients with CCD who were initiated on beta-blocker therapy for previous MI without a history of or current LVEF $\leq 50\%$, angina, arrhythmias, or uncontrolled hypertension, it may be reasonable to reassess the indication for long-term (> 1 year) use of beta-blocker therapy for reducing MACE. ⁹⁻¹⁵
3: No Benefit	B-NR	4. In patients with CCD without previous MI or LVEF $\leq 50\%$, the use of beta-blocker therapy is not beneficial in reducing MACE, in the absence of another primary indication for beta-blocker therapy. ^{†16-19}

Aim

To determine whether long-term oral beta-blocker treatment in patients with acute MI and preserved left ventricular function improves outcome

Outcome

- **Primary Endpoint**
 - Composite of death of any cause or new myocardial infarction (MI)
- **Secondary Endpoints**
 - All-cause death
 - Cardiovascular death
 - Myocardial infarction
 - Hospital admission due to atrial fibrillation
 - Hospital admission due to heart failure
- **Safety Outcomes**
 - Bradycardia, hypotension or Syncope
 - Hospitalization due to asthma or COPD
 - Stroke




Baseline characteristics

Characteristic	Beta-blockers (n=2508)	No Beta-blockers (n=2512)
Demography		
Median age (IQR) – year	65 (57-73)	65 (57-73)
Female sex, no (%)	563 (22.4)	568 (22.6)
Risk Factors		
Current smoker, no (%)	478 (19.4)	530 (21.3)
Hypertension, no (%)	1155 (46.1)	1163 (46.3)
Diabetes mellitus, no (%)	346 (13.8)	354 (14.1)
Prior cardiovascular disease		
Prior myocardial infarctions, no (%)	165 (6.6)	192 (7.7)
Prior PCI, no (%)	147 (5.9)	175 (7.0)
Prior CABG, no (%)	33 (1.3)	36 (1.4)
Prior Stroke, no (%)	52 (2.1)	67 (2.7)
Prior Heart failure, no (%)	13 (0.5)	22 (0.9)
Presentation characteristics		
Chest pain as main symptoms, no (%)	2421 (96.6)	2417 (96.2)
CPR before hospital, no (%)	10 (0.4)	11 (0.4)
Pulmonary rales, no (%)	29 (1.2)	42 (1.7)
Atrial fibrillation, no (%)	21 (0.8)	23 (0.9)
ST-elevation MI, no (%)	877 (35.0)	892 (35.5)
On oral beta-blocker treatment, no (%)	269 (10.9)	302 (12.2)

Baseline characteristics were similar between treatment groups

Primary outcome (all-cause death or MI)



Hypertension, the pressure is on!

BP goal <130/80 mmHg with GDMT*

1. GDMT

- ACE Inhibitors or ARB
- Thiazides
- DHP/NDHP CCBs
- Beta-blockers



<https://gfyca.com/totaltiredfinch>

*Guideline directed medical therapies depending on indications, e.g., ACE inhibitor or ARBs, beta-blockers

Differences in HTN categories

- JNC 7, JNC 8, and ACC/AHA 2017

2017 Guideline for the Prevention, Detection, Evaluation and Management of High Blood Pressure in Adults

BP Classification (JNC 7 and ACC/AHA Guidelines)

SBP		DBP	JNC 7	2017 ACC/AHA
<120	and	<80	Normal BP	Normal BP
120–129	and	<80	Prehypertension	Elevated BP
130–139	or	80–89	Prehypertension	Stage 1 hypertension
140–159	or	90–99	Stage 1 hypertension	Stage 2 hypertension
≥160	or	≥100	Stage 2 hypertension	Stage 2 hypertension

- Blood Pressure should be based on an average of ≥ 2 careful readings on ≥ 2 occasions
- Adults being treated with antihypertensive medication designated as having hypertension

HTN goals ACC/AHA 2017

Patient group	2017 ACC/AHA
General	<130/80 mm Hg
Older patients	<130 mm Hg [†]
Diabetes	<130/80 mm Hg
Chronic kidney disease	<130/80 mm Hg

*Includes patients with atherosclerotic cardiovascular disease (ASCVD) or an estimated 10-year risk $\geq 10\%$, as well as patients needing primary prevention or those with 10-year ASCVD risk $< 10\%$.

[†]General population ≥ 60 years of age. Treatment does not need to be adjusted in patients ≥ 60 years who may have lower systolic BP (eg, < 140 mm Hg) and are not experiencing adverse effects.

[‡]Ambulatory, community-dwelling, noninstitutionalized patients ≥ 65 years of age. Clinical judgment, patient preference, and a team-based approach to assess benefits and risks are reasonable for patients with a high burden of comorbidity and limited life expectancy.

Lifestyle first, foremost, and always



Consider discussing lifestyle modifications not as “work” you do to become healthy. Rather as doing enjoyable activities by yourself or with others that happen to help keep you feeling healthy.

Pharmacotherapeutics

- Initiation: what to start with? First-line and/or condition driven

Regardless of underlying conditions, **start with agents that have data for clinical outcomes benefits**, i.e., have clinical trial data demonstrating reduction of CVD events, CKD progression, etc.

Primary agents used in the treatment of hypertension include:

- **Thiazide diuretics** – (e.g., chlorthalidone, hydrochlorothiazide, indapamide, etc.)
- **ACE inhibitors*** – (e.g., enalapril, lisinopril, benazepril, etc.)*
- **ARBs*** – (e.g., candesartan, Olmesartan, irbesartan, losartan, etc.)
- **CCBs dihydropyridine** – (e.g., amlodipine, felodipine, nifedipine, etc.)
- **CCBs nondihydropyridine** – (e.g., diltiazem and verapamil)
- **B-blockers*** – (e.g., metoprolol succinate, carvedilol, bisoprolol)

*Class IB, preference towards ACEi/ARB and/or B-Blocker for HTN and/or MI/LV
Dysfunction

Specific diseases and populations

- BP goals (<130/<80) for all. Individuals and disease presence may differ.

- **Stable Ischemic Heart Disease** – GDMT ACEi/ARB +/- B-blockers
 - **Angina Pectoris** present DHP CCB thiazides, B-blockers
 - **Post-ACS**, LV dysfunction present B-blocker +/- ACEi/ARB; not present ACEi/ARB
e.g., lisinopril 5-10 mg/valsartan 80-160 mg, metoprolol succinate 25-50 mg, amlodipine 5-10 mg
- HFrEF – GDMT B-blockers, ACEi/ARB/ARNI, MRA. NDHP CCB NOT recommended.
- CKD – albuminuria (≥ 300 mg/day or ≥ 300 mg/g creatinine by first morning void) is present, ACEi, ARB if ACEi not tolerated. (**consider SGLT2i and ns-MRA**)
- DM – All first-line medications (e.g., thiazides, ACEi/ARB, DHP/NDHP CCBs) are reasonable.

Case

63-year-old woman presents for follow-up. She continues to have **stage 1 hypertension** and difficulty climbing stairs.

PMHx: Occasional headaches OB/GYN: Post-menopausal since early 20s EF 55%.

FMHx: Parents have passed. 2 brothers, 1 with DMII.

SocHx: Medical Technologist, working part-time. Married with 2 adult children. Does not follow any specific physical activity regimen.

MEDS: Clopidogrel 75 mg, rosuvastatin 40 mg, ezetimibe 10 mg, valsartan 160 mg

Vitals: **BP 138/80**, HR 80, SaO2 96%, T 98.9, Wt 155 Ht 5'5" BMI 25.8

LABS: TC 220, Trig 200, HDL 50, LDL 68. A1c 5.6%, Fasting Glucose 90

What are your recommendations?



Stage 1 - $\geq 130/\geq 80$, ASCVD $\geq 10\%$

1. Lifestyle modifications for health
2. Titrate BP medication: **Add amlodipine 5 mg (anti-anginal; titrate to 10 mg if needed) and consider increasing valsartan to 160 mg if needed**
3. Reiterate the importance of self-measurement and keeping a home BP journal
4. Reassess in 4-6 weeks in-person or by appropriate real-time communication (e.g., text, phone, or video visit)

In STATIN vs NO-STATIN, for every $\downarrow 38$ mg/dL LDL-C, there is a proportional $\downarrow 20\%$ in MACE in the 1st year.

After the 1st year, ongoing statin use leads to $\sim \downarrow 25\%$ in MACE in each subsequent year statin is used.

In more intensive STATIN vs less intensive STATIN, for each additional $\downarrow 19$ mg/dL LDL-C, an additional $\sim \downarrow 15\%$ in MACE





BOTTOMLINE:

High intensity statin = Each $\sim \downarrow 50$ mg/dL LDL-C, there is a $\sim \downarrow 28\%$ MACE

Secondary Prevention

STATIN...please.



MINIMUM 1st GOAL:

≥50% LDL-C Reduction from baseline.

High Intensity Statins (HIST)

- Atorvastatin 40, 80 mg
- Rosuvastatin 20, 40 mg

AHA/ACC 2018 2nd GOAL:

LDL-C <70 mg/dL OR non-HDL-C <100 mg/dL

Key TAKEAWAY in ASCVD:

1. Statin FIRST
2. Reduce LDL-C by >50% from baseline.
3. Add non-statins when LDL-C >70 or LDL >55
4. Check lipids 4-6 weeks after initiation or dose titration.

Updated AHA/ACC 2022:

ASCVD NOT at very high-risk LDL-C <70 mg/dL OR non-HDL-C <100 mg/dL

ASCVD at Very HIGH RISK, LDL-C <55 mg/dL OR non-HDL-C <85 mg/dL

Case

66-year-old man presents for follow-up. Returns for follow-up.

PMHx: Had an MI at 63, PCI w/2 stents to proximal LAD, EF 60%.
Type II Diabetes

FMHx: 1 brother with DMII

SocHx: Retired construction worker. Married with 1 adult child.
Former smoker, no EtOH. Walks daily for 40 minutes.

MEDS: Clopidogrel 75 mg, rosuvastatin 40 mg, ezetimibe 10 mg,
valsartan HCT 160/12.5 mg, dapagliflozin 10 mg, semaglutide 1.7
mg/weekly. SL NTG 0.4 mg PRN.

Vitals: BP 125/80, HR 80, SaO2 96%, T 98.9, Wt 155 Ht 5'5" BMI
25.8

LABS: TC 220, Trig 200, HDL 50, LDL 88. A1c 6.7%, UACR 40
mg/mmol. eGFR 92


What are your recommendations?

Very High-Risk ASCVD, LDL-C <55 mg/dL


1. Lifestyle modifications for health
2. Titrate lipid-lowering medication:
Add Repatha or Praluent (50% expected decrease, $88 - (88 \times 0.5) = 44$ mg/dL).
3. Reiterate the importance of self-measurement and keeping a home BP journal
4. Reassess in 4-6 weeks in-person or by appropriate real-time communication (e.g., text, phone, or video visit)

Diabetes Mellitus + CAD



DIABETES PREVALENCE



CVD PREVALENCE IN DIABETES



Diabetes Mellitus + CAD



Case

A 50-year-old woman presents for follow-up.

PMHx: DMII since age 30. HTN. MI at age 45; 3VCABG. EF 55%

FMHx: Mom with DMII. Dad with MI, age 70. 3 brothers, 2 with DMII.

SocHx: Director of Nursing. Married with 1 adult child. Lifetime nonsmoker, no EtOH. Five days/week gym class.

MEDS: Clopidogrel 75 mg, rosuvastatin 40 mg, valsartan 180 mg. SL NTG 0.4 mg PRN. Metformin 1000 mg 2 tabs QD, Lantus 30U daily, Insulin Aspart 15U with meals, glipizide 10 mg bid

Vitals: BP 140/80, HR 60, SaO2 96%, T 98.9, Wt. 200 Ht. 5'3"
BMI 35.4

LABS: TC 170, Trig 145, HDL 45, LDL 65. A1c 7.5%, Fasting Glucose 190 mg/dL

What are your recommendations?

Very High-Risk ASCVD, goal LDL-C <55 mg/dL

1. Lifestyle modifications for health
2. Add **ezetimibe 10 mg (20% expected to decrease, $65 - (65 \times 0.2) = 52$)**
3. **Add Amlodipine, Chlorthalidone, or Metoprolol Succinate**
4. **Add SGLT2i and/or GLP1ra and remove glipizide, reducing basal and short-acting insulin.**
5. Reassess labs in 4-6 weeks, with BP check, glucose journal (CGM?), by appropriate real-time communication (e.g., in-person text, phone, or video-visit)


ASCVD Sequelae

Death, nonfatal MI or stroke, PAD, etc.

- Angina – Optimal medical therapy or revascularization (PCI or CABG) + Optimal medical therapy
- Surveillance – Ankle Brachial Index, Carotid and/or abdominal ultrasound, stress tests

CV Mortality Risk by Albuminuria and eGFR Status in Patients With T2D^a

CV Mortality Risk



^aN=9795. ^bReference group has eGFR ≥ 90 and no albuminuria. ^cBaseline adjustment for age, sex, duration of diabetes, smoking, body mass index, systolic blood pressure, HbA1c, HDL-cholesterol, LDL-cholesterol, triacylglycerol, retinopathy, RAAS inhibition and treatment group. ^dIn mL/min/1.73 m². CV, cardiovascular; eGFR, estimated glomerular filtration rate; HDL, high-density lipoprotein; LDL, low-density lipoprotein; HbA1c, glycated hemoglobin; RAAS, renin-angiotensin-aldosterone system; T2D, type 2 diabetes.
Drury PL, et al. *Diabetologia*. 2011;54:32-43.

GFR and Albuminuria Are Predictive of CKD Progression

Prognosis of CKD by GFR and Albuminuria Categories: KDIGO 2012¹

				Persistent Albuminuria Categories Description and range		
				A1	A2	A3
				Normal to mildly increased	Moderately increased	Severely increased
				<30 mg/g <3 mg/mmol	30-300 mg/g 3-30 mg/mmol	>300 mg/g >30 mg/mmol
GFR Categories (mL/min/1.73 m ²) Description and range	G1	Normal or high	≥90			
	G2	Mildly decreased	60-89			
	G3a	Mildly to moderately decreased	45-59			
	G3b	Moderately to severely decreased	30-44			
	G4	Severely decreased	15-29			
	G5	Kidney failure	<15			

Figure reprinted with permission of The International Society of Nephrology: Kidney Disease Improving Global Outcomes Diabetes Work Group. *Kidney Int.* 2021;99:S1-S87.

- **A2** = *microalbuminuria*² (older classification system)
- **A3** = *macroalbuminuria or proteinuria*² (older classification system)

Risk of CKD Progression¹

- Low risk (if no other markers of kidney disease, no CKD)
- Moderately increased risk
- High risk
- Very high risk

CKD, chronic kidney disease; GFR, glomerular filtration rate; KDIGO, Kidney Disease Improving Global Outcomes.

MANAGEMENT

Slide credit: Dr. Mark Kraus


HTN in Chronic Kidney Disease; MRAs

- Nonsteroidal MRAs, Finerenone
- FIDELIO-DKD (CKD G3-4)

Outcome	Finerenone (N=2833) no. of patients with event (%)	Placebo (N=2841) no. of patients with event (%)	Finerenone (N=2833) no. of patients with event per 100 patient-yr	Placebo (N=2841) no. of patients with event per 100 patient-yr	Hazard Ratio (95% CI)	P Value
Primary composite outcome	504 (17.8)	600 (21.1)	7.59	9.08	0.82 (0.73–0.93)	0.001
Kidney failure	208 (7.3)	235 (8.3)	2.99	3.39	0.87 (0.72–1.05)	—
End-stage kidney disease	119 (4.2)	139 (4.9)	1.60	1.87	0.86 (0.67–1.10)	—
Sustained decrease in eGFR to <15 ml/min/1.73 m ²	167 (5.9)	199 (7.0)	2.40	2.87	0.82 (0.67–1.01)	—
Sustained decrease of ≥40% in eGFR from baseline	479 (16.9)	577 (20.3)	7.21	8.73	0.81 (0.72–0.92)	—
Death from renal causes	2 (<0.1)	2 (<0.1)	—	—	—	—
Key secondary composite outcome	367 (13.0)	420 (14.8)	5.11	5.92	0.86 (0.75–0.99)	0.03
Death from cardiovascular causes	128 (4.5)	150 (5.3)	1.69	1.99	0.86 (0.68–1.08)	—
Nonfatal myocardial infarction	70 (2.5)	87 (3.1)	0.94	1.17	0.80 (0.58–1.09)	—
Nonfatal stroke	90 (3.2)	87 (3.1)	1.21	1.18	1.03 (0.76–1.38)	—
Hospitalization for heart failure	139 (4.9)	162 (5.7)	1.89	2.21	0.86 (0.68–1.08)	—
Death from any cause	219 (7.7)	244 (8.6)	2.90	3.23	0.90 (0.75–1.07)	—
Hospitalization for any cause	1263 (44.6)	1321 (46.5)	22.56	23.87	0.95 (0.88–1.02)	—
Secondary composite kidney outcome	252 (8.9)	326 (11.5)	3.64	4.74	0.76 (0.65–0.90)	—
Sustained decrease of ≥57% in eGFR from baseline	167 (5.9)	245 (8.6)	2.41	3.54	0.68 (0.55–0.82)	—

Bakris et al., FIDELIO-DKD, NEJM, 2020.

A Urinary Albumin-to-Creatinine Ratio



No. of Patients

Finerenone	2831	2725	2582	1841	856
Placebo	2840	2726	2598	1825	834

Mean Change
from Baseline
(percent)

Finerenone	Ref.	-34.7	-41.3	-39.9	-29.3
Placebo	Ref.	-4.7	-3.0	-2.0	4.1

2023

SELECT TRIAL

M

Semaglutide and Cardiovascular
Outcomes in Obesity without Diabetes

multicenter, double-blind, randomized, placebo-controlled



Objective: to determine the association of subcutaneous semaglutide with cardiovascular (CV) events in a secondary prevention cohort of patients with overweight or obesity and prior CV disease (CVD) without diabetes mellitus (DM)

17,604
patients

Inclusion criteria: Age ≥ 45 years; BMI ≥ 27 kg/m²
Prior MI, stroke, or peripheral arterial disease (PAD)
with claudication and ankle-brachial index < 0.85 ,
prior revascularization, or amputation



Semaglutide
(n = 8803)

VS.



Placebo
(n = 8801)

PRIMARY OUTCOME

6.5

**composite of CV death, nonfatal MI,
and nonfatal stroke %**
HR 0.80, 95% CI 0.72-0.90, $p < 0.001$

8.0

SECONDARY OUTCOME

2.5

CV death %
HR 0.85, 95% CI 0.71-1.01, $p = 0.07$

3.0

3.4

CV death or HF hospitalization %
HR 0.82, 95% CI 0.71-0.96

4.1

Conclusion: In patients with preexisting cardiovascular disease and overweight or obesity but without diabetes, weekly subcutaneous semaglutide at a dose of 2.4 mg was superior to placebo in reducing the incidence of death from cardiovascular causes, nonfatal myocardial infarction, or nonfatal stroke at a mean follow-up of 39.8 months.

2024

FLOW TRIAL

M

Effects of Semaglutide on Chronic Kidney Disease
(CKD) in Patients with Type 2 Diabetes

A randomized, placebo controlled trial



Objective: To evaluate the safety and efficacy of semaglutide in slowing kidney function decline and reducing major adverse kidney events in patients with type 2 diabetes.

3533
Patients

Inclusion criteria: Age 18 years or older; diabetic, HbA1c $\leq 10\%$, renal impairment and undergoing treatment with maximum labelled or tolerated dose of ACE inhibitor or an ARB. **Exclusion criteria:** Congenital or autoimmune kidney disease, recent GLP-1 receptor agonist use or cardiovascular events, severe heart failure.



Semaglutide group
(n = 1767)

VS.



Placebo group
(n = 1766)

Primary Outcome

331

Major kidney disease events (N)
HR, 0.76; 95% CI, 0.66 to 0.88
($P = 0.0003$)

410

Secondary Outcomes

Slow

Decline in mean annual eGFR
(By 1.16 ml per minute per 1.73 m²)
($P < 0.001$)

Fast

49.6

Serious adverse events (%)

53.8

Conclusion: Semaglutide reduced the risk of clinically important kidney outcomes and death from cardiovascular causes in patients with type 2 diabetes and chronic kidney disease.

Common Questions

Cardiac evaluation for non-cardiac surgery (2022 ESC

<https://www.ahajournals.org/doi/10.1161/cir.0b013e3182447787>; 2014

AHA/ACC

<https://www.ahajournals.org/doi/full/10.1161/CIR.000000000000106>; Nice summary <https://www.acc.org/latest-in-cardiology/ten-points-to-remember/2022/09/01/13/18/2022-esc-guidelines-on-noncardiac-surgery-esc-2022>)

Return to work post cardiac bypass – work, severity, and patient dependent.

Intimacy and intercourse, 2012 AHA Scientific Statement

(<https://www.ahajournals.org/doi/10.1161/cir.0b013e3182447787>)

When to de-escalate therapies (age, cognitive, failure to thrive, terminal illnesses, etc.) – (Beers Criteria,

<https://geriatricscareonline.org/ProductAbstract/american-geriatrics-society-updated-beers-criteria/CL001/?param2=search>)